



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

able to maintain an apparently comfortable existence, unaided by man, under, to them, such strange environment. — J. A. ALLEN, *American Museum of Natural History, Central Park, New York City.*

Introduced Game Birds in Oregon and Idaho.—Several years since our consul to Japan sent over quite a number of Japanese Pheasants—exact species to me unknown. The birds were liberated below Portland, Oregon, on Sanvie's Island, and are known to have bred. While on duty at Vancouver Barracks I met a gentleman who informed me that he had seen, not long before (the summer of 1883), an old bird with several young near where the Pheasants were first liberated. In the autumn of 1883 a female was killed just below Vancouver. I saw this one, and the skin is now in the possession of Mr. John Jaggyat. The bird killed was one of a pair seen flying across the Columbia from the Oregon side to Washington Territory. The river at this point is nearly a mile wide, and the point of crossing was at least fifteen miles above the place of *planting*. By this it appears that the 'plant' has been established. The bird is a strong flier and bids fair to be a valuable acquisition. Oregon and Washington Territory have passed laws protecting these birds.

The Bob White (*Ortyx virginiana*) has been successfully introduced to the Boise Valley, Idaho. Three years since I found a covey on the west side of the Snake River, fifty miles below Boise City, where they were first liberated. I never saw coveys so large or numerous as I found them about Boise. Cover and food, as well as climate, are all favorable.—TIMOTHY E. WILCOX, *Assist. Surg., U. S. A.*

Fourth Addendum to List of Birds Ascertained to Occur within ten miles from Point de Monts, Province of Quebec, Canada; based chiefly upon the notes of Napoleon A. Comeau.—(For original List and previous Addenda see Bull. Nutt. Ornith. Club, Vol. VII, No. 4, Oct. 1882, pp. 233-242; Vol. VIII, No. 4, Oct. 1883, p. 244; and The Auk, Vol. I, No. 3, July 1884, p. 295; Vol. II, No. 1, Jan. 1885, p. 113.)

Mr. Comeau has recently sent me skins of the five following species:

174. *Dendroæca castanea*.—♂ shot at Godbout the last week in May, 1885.

175. *Dendroæca palmarum hypochrysea*.—♂ shot at Godbout, May 21, 1885.

176. *Cotile riparia*.—Shot at Godbout, June 8, 1885.

177. *Poœcetes gramineus*.—Shot at Godbout, April 24, 1885. Others were seen the same day and the next, and on May 10.

178. *Falco islandus*.—* A very handsome ♀ shot at Point de Monts, Jan. 7, 1885. Another was seen at Godbout, March 23, 1885.

179. *Bartramia longicauda*.—♂ shot at Godbout, May 7, 1885.

180. *Lobipes hyperboreus*.—Shot at Godbout, May 27, 1885.—C. HART MERRIAM, *Locust Grove, New York.*

* For authority for the name *Falco islandus* Brünn., see Stejneger in 'The Auk,' Vol. II, No. 2, April 1885, pp. 184-187.